

A neurophonetic perspective on articulation planning¹

Wolfram Ziegler

EKN, Institute of Phonetics, Ludwig-Maximilians-University, Munich, Germany
<https://doi.org/10.36505/ExLing-2020/11/0003/000418>

Abstract

This paper gives an overview of a model that predicts articulation ease for German phonological words on the basis of error data from patients with apraxia of speech (AOS). AOS is introduced as a clinical model of higher order motor processes for articulation. Word production accuracy in AOS is considered as a window into the structure of articulation plans as acquired through speech motor learning in childhood. The NLG model of apraxia of speech is explained. Applications in speech development and adult speech are outlined.

Keywords: articulation apraxia of speech

Neurophonetics

Neurophonetics is a research field that uses the knowledge and methodological tools developed in phonetics to study the neurological aspects of speaking. It covers investigations of the impact of brain dysfunctions on speech production in patients with neurologic conditions, but also of the neural substrates of speech in typical speakers.

Two different lines of neurophonetic research can be distinguished. First and foremost, phonetic thinking and methodologies are applied in the service of clinical neurology. An increasing number of studies is devoted to describing the speech patterns of patients with neurologic disorders, such as stroke, Parkinson's disease, or cerebellar ataxia. This research mainly aims at understanding the impact of neural dysfunction of specific aetiologies or localizations on the patients' speech characteristics. Applications of this research seek to establish physiologic, acoustic, or auditory-perceptual parameters that are sensitive and specific to neurologic speech impairments, with the ultimate goal of developing phonetically based clinical assessment tools (cf. Duffy 2019).

A second, less frequented line of neurophonetic research goes the opposite way, aiming to uncover general principles of speech production by investigating their breakdown due to dysfunctions of relevant brain networks. The study of disordered cognitive functions to learn more about the neural organization of "normal" cognition has long been a productive principle in cognitive neuropsychology, especially in neurolinguistics, though the "transparency" assumption underlying this principle has always been disputed. This assumption

claims that the impairment resulting from brain lesions is transparent for the structure of the underlying, unimpaired cognitive functions (Caramazza 1988). As a major precondition, this approach must be based on the solid foundations of a clinical model whose relationship to the cognitive function in question is sufficiently well established.

In this paper I will present a well-established clinical model of higher-order speech motor functions, *apraxia of speech* (AOS), and describe how investigations of this model may create evidence about the organization of typical speech. The disorder is considered to result from a dysfunction of the ability to plan the articulator movements required to produce the syllables, words, and sentences of the patient's native language.

A clinical model of articulation planning

Apraxia of speech occurs predominantly in patients who have suffered a left hemisphere stroke. Initially the patients are often completely unable to articulate, but within several hours or days their speech gradually recovers. Their articulation is usually slow, dysfluent, and effortful, and they produce errors such as substitutions, omissions and distortions of speech sounds, schwa intrusions in consonant clusters or cluster reductions, slowed or distorted transitions between speech sounds, and impaired coarticulation. They show visibly laborious groping movements of the articulators, with repeated attempts, false starts, and restarts in the initiation of utterances. The syndrome may manifest itself over a wide range of severity levels, from almost complete mutism to only mildly dysfluent speech with occasional articulation errors, and with different recovery dynamics across patients. For descriptions of the clinical pattern of AOS see e.g. Duffy (2019) or Ziegler (2008).

An important issue in the theoretical classification of this disorder is to differentiate it from other types of neurogenic sound production impairment. On the one hand, the symptoms of AOS are not explainable by "elementary" motor pathologies of the vocal tract muscles, such as paresis, ataxia, hypo- or hyperkinesia etc., which are summarized under the clinical term *dysarthria*. Unlike the dysarthrias, which are typically caused by bilateral brain lesions, AOS is a syndrome of the language-dominant hemisphere. The label "apraxia" historically relates to exactly this circumstance and characterizes the disorder as an impairment of "higher" motor functions. On the other hand, the syndrome is different from aphasic-phonological impairment, which is characterized by phoneme errors in essentially fluent speech, without any apparent signs of articulomotor involvement. Though these differential diagnostic considerations are far from trivial and have often been disputed, AOS is widely accepted as an autonomous clinical unit, and an enormous amount of research has been devoted to its clinical pattern and its neuro-anatomic substrate. For discussions see Ziegler, Aichert, and Staiger (2012).

Research interest in this condition has existed for almost 160 years. Broca's seminal case study of a man who lost "la faculté du langage articulé" after a lesion to the posterior part of the left inferior frontal gyrus (Broca, 1861) is generally acknowledged as the birth of systematic clinical brain-behaviour research. Broca pointed out, in the terminology of his time, that the patient was unable to articulate, although he had no obvious motor restrictions of the tongue and lips and no generalized language or cognitive impairment. He allocated the ability to articulate to the left frontal cortical region that bears his name until today, and characterized the speech impairment as a "loss of the memory of the procedures required for the articulation of words" (Broca 1861; p. 333).

Since Broca's time, the neuro-anatomic basis and the functional characterization of this condition have been discussed extensively, - probably more than any other neurologic speech or language dysfunction. Over decades, evidence has accumulated that lesions to left posterior inferior frontal cortex including the opercular part of Broca's area and the adjacent pre-motor and motor cortex, as well as subjacent anterior insular cortex, are responsible for the development of AOS. Moreover, in more recent years this region has been identified in numerous imaging studies of non-brain-damaged speakers as a higher-order speech motor centre. The dysfunction resulting from lesions to this cortical area has been characterized in varying terms as "apraxia of the language muscles", "phonetic disintegration of speech", a "programming deficit", a breakdown of the "functional coalitions" of articulation, or, more recently, a disorder of "phonetic planning" or "speech motor planning" (for a historical review see Ziegler et al. 2012).

Implied in this thinking is that the pathomechanism of AOS disrupts the language-specific motor patterns for the production of syllables, words and sentences as acquired during speech development. We have recently proposed a model that delineates how speech motor learning in childhood is mediated by subcortical structures and leads to an accumulation of a "knowledge base for articulation" in the left frontal cortex of the adult brain (Ziegler & Ackermann 2017). An obvious assumption, based on a wealth of knowledge about the learning-dependent plasticity of the human brain, is that speech motor patterns that are more strongly integrated through speech learning are represented more redundantly within the functional network of this brain region. By implication, such patterns are less vulnerable to a loss of cortical tissue in this area. Conversely, articulatory patterns that are less typical of the speaker's native language, or less cohesive, have less redundant neural representations and are therefore more vulnerable to cortical damage. For a more detailed explanation of this argument see Ziegler, Lehner, Pfab & Aichert (2020).

As a conclusion, the speech patterns of patients with AOS provide a window into the make-up of the acquired implicit knowledge about how the words and sentences of their language are articulated. The argument is similar to that

brought up in studies of speech errors in healthy speakers (e.g., Pouplier & Hardcastle 2005), but with the difference that a much larger corpus of data can be acquired from patients with AOS under relatively natural speaking conditions, and that the errors made by patients with AOS can be allocated to a rather circumscribed functional and neuro-anatomic source, i.e., a dysfunction of the acquired articulation planning processes located in the left posterior inferior frontal lobe.

Accuracy of apraxic word production as a yardstick of “articulatory ease”

For the reasons mentioned above, the errors made by patients with AOS reveal what is easy and what is difficult to articulate for adult native speakers of a given language. “Ease of articulation” is a disputed concept, at least for adult speech, because the highly overlearned nature of the articulation patterns of our native language entails that everything is equally easy for us to say (Ladefoged 1990). In the sense used here, a word is easy to pronounce if patients with apraxia of speech have relatively few problems producing it. More specifically, patients with only mild impairment may mostly produce it correctly, and only those with severe AOS make errors on it. Conversely, difficult phonetic patterns are those that provoke errors even in patients with mild AOS. Following the transparency assumption explained above, the susceptibility of words to apraxic failure mirrors the redundancy of the representation of its motor components in left inferior frontal cortex, which, in turn, is considered the neural substrate of the degree to which language-specific articulatory patterns were stabilized through speech motor learning during childhood.

In many of the earlier studies of AOS, error rates were related to phonemes, mostly with the finding that consonants are less error prone than vowels, plosives and nasals less than fricatives or affricates, voiceless obstruents less than voiced obstruents, etc. Other findings were related to syllable structure, e.g., that coda consonant errors are less frequent than onset errors, or simplex syllables are less vulnerable than complex syllables. Finally, at the supra-syllabic level a rather common finding was that the likelihood of an error increases with the number of syllables in a word. More recently, we could also demonstrate an effect of lexical stress, showing that in German AOS patients, trochaic words were “easier” to pronounce than iambic words (Aichert, Späth & Ziegler 2016). An overview of these findings including the corresponding references is listed in Table 1 of Ziegler et al. (2020).

The NLG-model

Considering that the hierarchies of articulatory planning requirements sketched above extend across all levels of the phonological architecture of words, an approach that integrates these levels is necessary to account for the interactions

between them. The Nonlinear Gestural (NLG) model of AOS, which is based on ideas from Articulatory Phonology (e.g., Goldstein, Pouplier, Chen, Saltzman, & Byrd 2007) considers articulatory gestures as the basic elements of articulation planning and postulates that these units are successively integrated to form syllable structure components, syllables, metrical feet, and phonological words. The model has been described in more detail elsewhere (Ziegler & Aichert 2015; Ziegler, Aichert, & Staiger 2017; Ziegler et al. 2020). In brief, it postulates that in a speaker with AOS of a given degree of severity, the likelihood that an articulatory gesture is planned accurately assumes a certain value $p \in]0,1[$. The more severe the disorder, the more likely is that the production of a gesture fails, hence the smaller is p . By combinatorial laws, the probability that a combination of two gestures is correct drops to p^2 , but only under the unlikely assumption that they are independent. More generally, the likelihood that two combined gestures are accurate is expressed as $p^2 * c$, with an unknown coefficient c indicating whether the probability of accurate production increases ($c > 1$) or decreases ($c < 1$). An increase would point at some inherent bonding of the two gestures, either by biomechanical conditions or through a learned gestural integration, which makes them less vulnerable because they do not count as two independent occasions to fail. Conversely, a decrease would suggest that putting the two gestures together requires additional effort, which makes them more vulnerable than predicted from purely combinatorial considerations. Hence, the coefficients in the model represent the “glue” between the gestural components of syllables and words. Depending on the structural relationship of a gesture with other gestures in a word, different coefficients are inserted to represent the specific degree of integration of the respective gesture within the word’s gestural architecture.

Figure 1 illustrates, as an example, the gestural score of the three-syllabic word *Trompete* (engl. *trumpet*). The figure shows a variety of inter-gestural relationships implemented in the NLG-model, such as synchrony of a vocal tract gesture with a glottal (e.g., /t/) or velar gesture (e.g., /m/), consonantal gestures in pre- or post-vocalic positions, gestures clustered within a pre-or post-vocalic position, gestures in the tail of a metrical foot (/tə/) attached to those in the head (/pe:/), and those of an extrametrical syllable to the gestures within a metrical foot. Computationally, each word is modelled by a nonlinear, multiplicative term representing each gesture’s base probability of correct production and the coefficients coding for the gesture’s bonding strengths within the word’s gestural score, as displayed in figure 1.




Figure 1: Gestural score of the 3-syllabic German word Trompete (with SAMPA transcript). Each articulatory gesture is represented by a trapezoid. Consonant gestures are arranged on different layers depending on the involved constriction organ (LIP: lips, TT: tongue tip, TB: tongue back, VEL: velar aperture, GLO: glottal aperture). Closed constrictions are represented by filled trapezoids, the “critical” constriction type of the /ʁ/ gesture by a patchy trapezoid. Vowel gestures (in yellow) are allocated to a separate layer representing the gross vocal tract deformations of vowel articulation, including jaw, tongue body and lip rounding movements. Dashed vertical lines indicate syllable boundaries, the tree structure characterizes the word’s stress pattern (thick lines indicating prominence).

Model coefficient estimates were calculated and validated in nonlinear regression analyses using corpora of 96 and 136 words of different lengths and complexities, with 120 and 66 renditions per word, respectively, by patients with AOS (Ziegler & Aichert 2015; Ziegler et al. 2020).

Model shape

The sizes of the model coefficients indicate where particularly strong or particularly weak cohesions exist within the gestural patterns of words. The results largely corroborate earlier findings about the factors influencing accuracy of apraxic speech, but unlike earlier investigations of isolated factors, the data present an integrative view that takes account of the interactions between the hierarchical levels of the model.

Here is a selection of findings: onset gestures are more demanding than coda gestures, glottal and velar aperture gestures are highly cohesive with their synchronous tongue- or lip gestures, cluster gestures are independent, gestures within the tail-position of a metrical foot are inexpensive, whereas left-branching expansions of metrical feet are expensive (Ziegler et al. 2020). A transfer of these results to the motor organization of typical, neurologically

healthy speech is only possible on a relative scale. For example, the finding that the coefficient representing the cohesive ties of two cluster gestures is close to 1 suggests an independent recruitment of both gestures, i.e., no integration, in patients with AOS. This does not preclude that gestures in consonant clusters are organized as cohesive units in healthy speakers (Hoole & Pouplier 2015). However, the ties between a tongue back closure and glottal aperture gesture in the production of /k/, with a coefficient considerably greater than 1, are stronger than those between the same tongue back gesture and a tongue tip closure in the cluster /kn/.

Applications

Beyond the clinical applications of the NLG model (Ziegler et al. 2020), this research offers options for investigations of typical speech in adults or in children. A web application for calculating an NLG-score for each word or nonword that meets German phonotactic constraints can be accessed at <https://neurophonetik.de/sprechapraxie-gestenmodell>. This tool allows for the computation of NLG scores for large lexical databases, (e.g., SUBTLEX-DE, a German lexicon containing word frequency data, <https://neurophonetik.de/subtlex-np>), with options to investigate the relationships between articulatory ease on the one hand and lexical frequency, lexical familiarity, neighbourhood density, or age of acquisition on the other (e.g., Lehner & Ziegler, accepted). Furthermore, corpora of child word forms can be examined for their relationships with their corresponding adult forms, to investigate hypotheses concerning the role of articulation in infant word learning. Examples of such data will be included in the oral presentation.

Acknowledgement

This research was supported by grants from the German Research Council (DFG, Zi 469 / 6-1/8-1/14-1,2). I am grateful to Ingrid Aichert and Anja Staiger for their long-standing valuable contributions to this work and to the many patients who have agreed to participate in our research.

Notes

1. An expanded version of this paper was submitted to Laboratory Phonology An expanded version of this paper was submitted to Laboratory Phonology.
2. See www.neurophonetik.de/gesten-koeffizientenrechner.

References

- Aichert, I., Späth, M., Ziegler, W. 2016. The role of metrical information in apraxia of speech. Perceptual and acoustic analyses of word stress. *Neuropsychologia* 82, 171-178.

- Broca, P. 1861. Remarques sur le siège de la faculté du langage articulé; suivies d'une observation d'aphémie (perte de la parole). *Bulletins et memoires de la Société Anatomique de Paris*, XXXVI 330-357.
- Caramazza, A. 1988. Some aspects of language processing revealed through the analysis of acquired aphasia: The Lexical System. *Ann Rev Neurosci* 11, 395-421.
- Duffy, J. 2019. *Motor Speech Disorders: Substrates, Differential Diagnosis, and Management*: Elsevier Health Sciences.
- Goldstein, L., Pouplier, M., Chen, L., Saltzman, E., Byrd, D. 2007. Dynamic action units slip in speech production errors. *Cognition* 103, 386-412.
- Hoole, P., Pouplier, M. 2015. Interarticulatory Coordination. *Speech Sounds*. In M. A. Redford (ed.), *The Handbook of Speech Production*, 133-157. Chichester: Wiley.
- Ladefoged, P. 1990. Some reflections on the IPA. *J Phonet* 18, 335-346.
- Lehner, K., Ziegler, W. accepted. The impact of lexical and articulatory factors in the automatic selection of test materials for a web-based assessment of intelligibility in dysarthria. *J Speech Lang Hear Res*
- Pouplier, M., Hardcastle, W. 2005. A re-evaluation of the nature of speech errors in normal and disordered speakers. *Phonetica* 622, 227-243.
- Ziegler, W. 2008. Apraxia of speech. In G. Goldenberg M. Miller (eds.), *Handbook of Clinical Neurology*, 269-285. London: Elsevier.
- Ziegler, W., Ackermann, H. 2017. Subcortical Contributions to Motor Speech: Phylogenetic, Developmental, Clinical. *Trends Neurosci* 40, 458-468.
- Ziegler, W., Aichert, I. 2015. How much is a word? Predicting ease of articulation planning from apraxic speech error patterns. *Cortex*, 69, 24-39.
- Ziegler, W., Aichert, I., Staiger, A. 2012. Apraxia of speech: Concepts and controversies. *Jf Speech Lang Hear Res* 55(5), S1485-S1501.
- Ziegler, W., Aichert, I., Staiger, A. 2017. When words don't come easily: A latent trait analysis of impaired speech motor planning in patients with apraxia of speech. *J Phonet* 64, 145-155.
- Ziegler, W., Lehner, K., Pfab, J., Aichert, I. 2020. The nonlinear gestural model of speech apraxia: clinical implications and applications. *Aphasiology*, 1-23. doi:10.1080/02687038.2020.1727839